

DRUG FACTS

Active ingredient(s)

Purpose

Use(s)

temporarily relieves these symptoms associated with hay fever or other upper respiratory allergies, and the common cold:

- runny nose
- itchy, watery eyes
- itching of the nose or throat
- sneezing
- nasal congestion
- sinus pressure
- headache
- minor aches and pains
- fever

Warnings

Do not use

- in children under 12 years of age
- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- problems or serious side effects from taking pain relievers or fever reducers
- stomach problems that last or come back, such as heartburn, upset stomach, or stomach pain
- ulcers
- bleeding problems
- high blood pressure
- heart or kidney disease
- asthma
- thyroid disease
- diabetes
- glaucoma
- trouble urinating due to an enlarged prostate gland
- taken a diuretic
- reached age 60 or older

Ask a doctor or pharmacist before use if you are

- taking any other drug containing a NSAID (prescription or nonprescription)
- taking a blood thinning (anticoagulant) or steroid drug
- under a doctor's care for any serious condition
- taking sedatives or tranquilizers
- taking any other product that contains pseudoephedrine, chlorpheniramine or any other nasal decongestant or antihistamine
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding.
- fever gets worse or lasts more than 3 days
- nasal congestion lasts for more than 7 days
- stomach pain or upset gets worse or lasts
- redness or swelling is present in the painful area
- you get nervous, dizzy, or sleepless
- symptoms continue or get worse
- any new symptoms appear

If pregnant or breast-feeding

Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

Other information

Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing an NSAID [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

- **do not take more than directed**
- **the smallest effective dose should be used**
- do not take longer than 7 days, unless directed by a doctor (see WARNINGS)
- adults: take 1 caplet every 4-6 hours while symptoms persist.
- do not take more than 6 caplets in any 24-hour period, unless directed by a doctor
- children under 12 years of age: do not use
- read all warnings and directions before use. Keep carton.
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients

carnauba wax, colloidal silicon dioxide, corn starch, croscarmellose sodium, FD&C red no. 40 aluminum lake, FD&C yellow no. 6 aluminum lake, glyceryl behenate, hypromellose, microcrystalline cellulose, pharmaceutical ink, polydextrose, polyethylene glycol, pregelatinized starch, propylene glycol, silicon dioxide, titanium dioxide